


**COVID-19 Testing Algorithm for Antepartum Obstetric Patients Who Require Admission or Procedure**

**Screen for COVID-19**

In the last 30 days, have you been in contact with anyone who was confirmed or suspected to have COVID-19, including outside the US?

**OR**, do you have any of the following: Fever  $\geq 100.0^{\circ}\text{F}$  or  $37.8^{\circ}\text{C}$ , cough, shortness of breath, chills, shaking chills, fatigue, muscle pain, headache, sore throat, new loss of taste or smell, congestion or runny nose, nausea or vomiting, diarrhea

**OR**, have you been advised to stay home by a medical professional due to illness that might be COVID-19?



**\* A person is cleared from COVID-19 infection when 14 days from onset of symptoms AND 3 or more days from resolution of fever, whichever is longer.**

**Additional Guidance Regarding COVID-19 Nasopharyngeal Swabbing of the Obstetrical or postpartum patient**

- Patient to wear surgical mask over mouth during swab collection to help minimize any spread of droplets
- Scheduled **CD or ECV** patient will have swab obtained the day PRIOR to scheduled **CD or ECV**
- For the asymptomatic SVD patient, the preferred swabbing location is in her labor and birth or MSCU room with the door closed. If unavailable, next best location is their triage room with the door closed.
- For the asymptomatic preoperative CD or planned CD patient, the preferred swabbing location is in their PACU room with the door closed.
- For the asymptomatic antepartum or postpartum admission, the preferred swabbing location in their hospital room with the door closed.

1.

	COVID-19 monitoring
	Inpatient COVID-19 Test for Clearance or Disposition WITHOUT Symptoms
	Inpatient COVID-19 Test PUI or WITH Symptoms
	LORAZEPAM (ATIVAN) CONTINUOUS INFUSION (ADULT ICU) (aka Y RX COVID19 LORAZEPAM INF SHORTAGE)

2.

✔ Accept

**Inpatient COVID-19 Test for Clearance or Disposition WITHOUT Symptoms**

**Patients do not need to be moved to a new room**

**DO NOT modify (or add) isolation precautions based upon this order**

**\*\*Guidance on testing for discharges to SNF\*\***  
For COVID positive patients:  
 - Do NOT retest until 14 days **after** the initial test (not a prerequisite for discharge)  
 - Clarify if there is a SNF willing to accept the patient without retesting  
 - If patient remains hospitalized for ≥14 days **and** symptoms resolved, order two repeat tests 24 hours apart

For non- COVID patients requiring SNF:  
 - Order within 24 hours of planned discharge

✔ Accept ✖ Cancel

**✔ COVID-19 Clearance or Disposition (\$\$\$\$\$)**

Frequency:  Once STAT Tomorrow AM

Starting:  Today Tomorrow At:

First Occurrence: **Today 1052**

Scheduled Times ⤴

09/30/20 1052

Specimen Type:

Specimen Src:

Lab Specific Advisory Note:  DO NOT PLACE ORDERS IN COMMENT FIELD

⚠ 2 hour test extremely limited Lab may need to run on less rapid platform

⚠ Active Labor or OR/Procedure Within 4 Hours Yes No

Comments: ✎ Patients do not need to be moved to a new room. DO NOT modify (or add) isolation precautions based upon this order.

Phase of Care:

Add-on: No add-on specimen found

✔ Accept ✖ Cancel

✔ Accept

⚠ Next Required

